

PEORIA AREA EMS SYSTEM
PREHOSPITAL CARE MANUAL

**Heat-Related Emergencies
Protocol**

Injury and illness from heat exposure varies depending on the *manner* of exposure (sun, humidity, exertion) and the *amount* of exposure (time, temperature & ambient air).

Heat exposure emergencies range from localized cramping to severe hyperthermia (heat stroke) with unresponsiveness and unconsciousness. The patient's health, predisposing factors and medications may increase the likelihood of heat-related illness and injury. The primary goal in the treatment of the patient at risk for hyperthermia is to cool the patient and restore body fluids.

First Responder Care

First Responder Care should be focused on assessing the situation and initiating routine patient care to assure that the patient has a patent airway, is breathing and has a perfusing pulse as well as beginning treatment for shock.

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. Move the patient to a cool environment. Remove clothing as necessary to make the patient comfortable. Cold packs may be utilized for the neck (posterior), armpits, groin and along the thorax. Do not cool the patient to a temperature that will cause them to shiver.
3. **Oxygen:** 15 L/min via non-rebreather mask or 6 L/min via nasal cannula if the patient cannot tolerate a mask. Be prepared to support the patient's respirations with BVM if necessary.

BLS Care

BLS Care should be directed at conducting a thorough patient assessment, initiating routine patient care to assure that the patient has a patent airway, is breathing and has a perfusing pulse as well as beginning treatment for shock and preparing the patient for or providing transport.

1. Render initial care in accordance with the *Routine Patient Care Protocol*.

PEORIA AREA EMS SYSTEM
PREHOSPITAL CARE MANUAL

**Heat-Related Emergencies
Protocol**

BLS Care (continued)

2. Move the patient to a cool environment. Remove clothing as necessary to make the patient comfortable. Cold packs may be utilized for the neck (posterior), armpits, groin and along the thorax. Do not cool the patient to a temperature that will cause them to shiver.
3. **Oxygen:** 15 L/min via non-rebreather mask or 6 L/min via nasal cannula if the patient cannot tolerate a mask. Be prepared to support the patient's respirations with BVM if necessary.
4. Treat other symptoms per the appropriate protocol.
5. Initiate ALS intercept if needed and transport as soon as possible.

ILS Care

ILS Care should be directed at continuing or establishing care, conducting a thorough patient assessment, stabilizing the patient's perfusion and preparing for or providing patient transport.

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. Move the patient to a cool environment. Remove clothing as necessary to make the patient comfortable. Cold packs may be utilized for the neck (posterior), armpits, groin and along the thorax. Do not cool the patient to a temperature that will cause them to shiver.
3. **Oxygen:** 15 L/min via non-rebreather mask or 6 L/min via nasal cannula if the patient cannot tolerate a mask. Be prepared to support the patient's respirations with BVM if necessary.
4. **IV Fluid Therapy:** 500mL fluid bolus if the patient is hypotensive to achieve a systolic BP of at least 100mmHg.
5. Treat other symptoms per the appropriate protocol.
6. Initiate ALS intercept if needed and transport as soon as possible.

PEORIA AREA EMS SYSTEM
PREHOSPITAL CARE MANUAL

**Heat-Related Emergencies
Protocol**

ALS Care

ALS Care should be directed at continuing or establishing care, conducting a thorough patient assessment, stabilizing the patient's perfusion and preparing for or providing patient transport.

1. Render initial care in accordance with the *Routine Patient Care Protocol*.
2. Move the patient to a cool environment. Remove clothing as necessary to make the patient comfortable. Cold packs may be utilized for the neck (posterior), armpits, groin and along the thorax. Do not cool the patient to a temperature that will cause them to shiver.
3. **Oxygen:** 15 L/min via non-rebreather mask or 6 L/min via nasal cannula if the patient cannot tolerate a mask. Be prepared to support the patient's respirations with BVM if necessary.
4. **IV Fluid Therapy:** 500mL fluid bolus if the patient is hypotensive to achieve a systolic BP of at least 100mmHg.
5. Treat other symptoms per the appropriate protocol.
6. Transport as soon as possible.

PEORIA AREA EMS SYSTEM
PREHOSPITAL CARE MANUAL

**Heat-Related Emergencies
Protocol**

Heat Disorders

Heat (Muscle) Cramps – Heat cramps are muscle cramps caused by overexertion and dehydration in the presence of high temperatures. Signs & symptoms include: *Normal or slightly elevated body temperature; generalized weakness; dizziness; warm, moist skin and cramps in the fingers, arms, legs or abdominal muscles.*

Heat Exhaustion – Heat exhaustion is an acute reaction to heat exposure and the most common heat-related illness a prehospital provider will encounter. Signs & symptoms include: *Increased body temperature; generalized weakness; cool, diaphoretic skin; rapid, shallow breathing; weak pulse; diarrhea; anxiety; headache and possible loss of consciousness .*

Heatstroke – Heatstroke occurs when the body's hypothalamic temperature regulation is lost. Cell death and damage to the brain, liver and kidneys can occur. Signs & symptoms include: *Cessation of sweating; very high core body temperature; hot, usually dry skin; deep, rapid, shallow respirations (which later slow); rapid, full pulse (which later slows); hypotension; confusion, disorientation or unconsciousness and possible seizures.*

Fever (Pyrexia) – A fever is the elevation of the body temperature above the normal temperature for that person (~ 98.6° F +/- 2 degrees). Fever is sometimes difficult to differentiate from heatstroke; however, there is usually a history of infection or illness with a fever.