

**PEORIA AREA EMS SYSTEM
PEDIATRIC PREHOSPITAL CARE MANUAL**

**Pediatric Assessment Process
and Management**

Pediatric Age Definitions & Assessment Considerations

Neonate (0-1 Month):

- Utilization of APGAR Scoring is helpful in assessing the neonate patient.

Infant (1-12 Months):

- Approach the infant slowly and calmly. Fast motion and loud noises may startle or agitate the infant.
- Use warm hands and assessment tools.
- Avoid doing anything potentially painful or distressing until after the assessment is completed.
- Have the caregiver assist in care – this is less threatening to the infant.
- Children over six (6) months of age are usually best examined in the arms of a parent. “Stranger anxiety” may be present and could eliminate other assessment options.
- If needed, calm the infant with a pacifier, blanket or favorite toy.

Toddler (1-3 Years):

- Approach the toddler slowly. Keep physical contact at a minimum until he/she feels familiar with you.
- Perform the assessment at the level of the toddler by sitting or squatting next to them and **allow the toddler to remain in the caregiver’s lap** whenever possible.
- Assessment should be **toe to head**. This is less threatening to the toddler.
- Give limited choices such as “Do you want me to listen to your chest or feel your wrist first?”

PEORIA AREA EMS SYSTEM
PEDIATRIC PREHOSPITAL CARE MANUAL

**Pediatric Assessment Process
and Management**

Pediatric Age Definitions & Assessment Considerations

Toddler (1-3 Years) {Continued}:

- Use simple, concrete terms and continually reassure the toddler.
- Do not expect the toddler to sit still and cooperate – be flexible.

Preschooler (3-5 Years):

- A preschool aged child is a “magical thinker.” Concrete concepts must be described in short, simple terms.
- A preschooler is often very cooperative during the assessment process and may be able to provide a history.
- Questions should be simple and direct.
- Allow the child to handle equipment.
- Use distractions.
- **Do not lie to the child. If the procedure is going to hurt, tell them.**
- Set limits on behavior (*e.g.* “You can cry or scream, but don’t bite or kick.”).
- Focus on one thing at a time.
- Play games with immobilized preschoolers to distract him/her and prevent them from squirming.

**PEORIA AREA EMS SYSTEM
PEDIATRIC PREHOSPITAL CARE MANUAL**

**Pediatric Assessment Process
and Management**

Pediatric Age Definitions & Assessment Considerations

School Age (5-13 Years):

- The school aged child is usually cooperative and can be the primary sources for the patient history.
- Explain all procedures simply and completely and respect the patient's modesty.
- Substance abuse issues may be present in this age group and should be considered during the care of altered level of consciousness cases.
- Children at this age are afraid of losing control, so let him/her be involved in the care. However, do not negotiate patient care unless the child really has a choice.
- Reassure the child that being ill or injured is not a punishment and praise them for cooperating.

Adolescent (13-16 Years):

- **The adolescent is more of an adult than a child and should be treated as such.** Depending on the nature of the problem, an accurate history may not be possible with parents observing. It may be necessary to separate the parent and child during the assessment.
- Regardless of who is present, respect the patient's modesty. Avoid exposing the adolescent unnecessarily.
- Explain what you are doing and *why* you are doing it!
- Show respect – speak to the adolescent directly. Do not turn to the caregiver for the initial information.