

## Continuing Education Quiz (1.0 hour CEU)

Name: \_\_\_\_\_

Address: \_\_\_\_\_

City: \_\_\_\_\_ State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

Email: \_\_\_\_\_

### Factors Associated with Cervical Spine Injury in Children After Blunt Trauma

- 1) According to this article, cervical spine injury occurs in up to 10% of children presenting for trauma evaluation

- True  
 False

- 2) Which of the following is **NOT** a common and known adverse effect of cervical spine immobilization and radiographic assessment

- Pain  
 Bleeding  
 Pressure wounds  
 Exposure to ionizing radiation

- 3) Which of the following best describes what type of study this was?

- Double blinded, placebo controlled  
 Retrospective, chart review  
 Prospective, case control  
 Retrospective, case control

- 4) In this article, when defining altered mental status, which of the following was **NOT** included?

- Glasgow Coma Scale (GCS) < 15  
 Evidence of intoxication  
 AVPU < A  
 Speaking a full sentence

- 5) A 12 year old who was injured after diving into a pool is considered a high-risk mechanism.

- True  
 False

- 6) In this proposed model, posterior neck tenderness was identified as an important part of the criteria.

- True  
 False

7) This study and the 8 identified criteria will require further investigation prior to use in a clinical setting.

- True
- False